

ssresp.eig Linear algebraic eigenproblem

1 The linear algebraic eigenproblem can be simply stated. For $n \times n$ real matrix A , $n \times 1$ complex vector \mathbf{m} , and $\lambda \in \mathbb{C}$, \mathbf{m} is defined as an eigenvector of A if and only if it is nonzero and

$$A\mathbf{m} = \lambda\mathbf{m} \quad (1)$$

for some λ , which is called the corresponding eigenvalue. That is, \mathbf{m} is an eigenvector of A if its linear transformation by A is equivalent to its scaling; i.e. an eigenvector of A is a vector of which A changes the length, but not the direction.

2 Since a matrix can have several eigenvectors and corresponding eigenvalues, we typically index them with a subscript; e.g. \mathbf{m}_i pairs with λ_i .

Solving for eigenvalues

Eq. 1 can be rearranged:

$$(\lambda I - A)\mathbf{m} = \mathbf{0}. \quad (2)$$

For a nontrivial solution for \mathbf{m} ,

$$\det(\lambda I - A) = 0, \quad (3)$$

which has as its left-hand-side a polynomial in λ and is called the characteristic equation. We define eigenvalues to be the roots of the characteristic equation.

Box ssresp.1 eigenvalues and roots of the characteristic equation

If A is taken to be the linear state-space representation A , and the state-space model is converted to an input-output differential equation, the resulting ODE's "characteristic equation" would be identical to this matrix characteristic equation. Therefore, everything we

eigenproblem

eigenvector

eigenvalue

characteristic equation
eigenvalues

already understand about the roots of the “characteristic equation” of an i/o ODE—especially that they govern the transient response and stability of a system—holds for a system’s A -matrix eigenvalues.

3 Here we consider only the case of n distinct eigenvalues. For eigenvalues of (algebraic) multiplicity greater than one (i.e. repeated roots), see the discussion of [Appendix adv.eig](#).

Solving for eigenvectors

4 Each eigenvalue λ_i has a corresponding eigenvector \mathbf{m}_i . Substituting each λ_i into [Eq. 2](#), one can solve for a corresponding eigenvector. It’s important to note that an eigenvector is unique within a scaling factor. That is, if \mathbf{m}_i is an eigenvector corresponding to λ_i , so is $3\mathbf{m}_i$.³

3. Also of note is that λ_i and \mathbf{m}_i can be complex.

Example [ssresp.eig](#) – 1

Let

$$A = \begin{bmatrix} 2 & -4 \\ -1 & -1 \end{bmatrix}.$$

• Find the eigenvalues and eigenvectors of A .

re: eigenproblem for a 2×2 matrix



5 Several computational software packages can easily solve for eigenvalues and eigenvectors. See [Lec. ssresp.eigcomp](#) for instruction for doing so in Matlab and Python.